



CLINICAL GUIDELINE	
Cardiac: Blalock-Taussig Shunt (BT Shunt) Management Following Insertion	
Scope (Staff):	Nursing and Medical Staff
Scope (Area):	NICU KEMH, NICU PCH, NETS WA
Child Safe Organisation Statement of Commitment	
<p>The Child and Adolescent Health Service (CAHS) commits to being a child safe organisation by meeting the National Child Safe Principles and National Child Safe Standards. This is a commitment to a strong culture supported by robust policies and procedures to ensure the safety and wellbeing of children at CAHS.</p>	

This document should be read in conjunction with this [DISCLAIMER](#)

Contents

Surgical Procedure	2
Routine Post-Op Management Following Neonatal BT Shunt Insertion.....	2
Post-Op BT Shunt Complications.....	3
Low O2 Saturations (< 70%).....	4
High O2 Saturations (> 85% in Air)	4
Prior to Discharge	5
References.....	2

Aim

This guideline focuses on the specifics of the post-op management and complications of a neonate following the insertion of a Blalock-Taussig (BT) shunt. For routine post-op cardiac care, general complications following cardiac surgery and for general information on tricuspid atresia/ pulmonary atresia and Tetralogy of Fallot, and their pre-op management please refer to the following guidelines


- [Cardiac: Routine Post-Operative Care](#)
- [Cardiac: Complications Management Following Surgery](#)

A BT shunt is a systemic to pulmonary shunt which is a palliative procedure usually performed in neonates to establish a reliable pulmonary blood flow in patients with obstructed pulmonary blood flow e.g. tricuspid atresia, pulmonary atresia and severe Tetralogy of Fallot. The pulmonary blood flow will have been maintained pre-operatively by maintaining ductal patency with [Prostaglandin E1 \(PGE1\)](#). The definitive operation/s is/are performed later when the child is larger.

Surgical Procedure

The Classical BT Shunt (CBTS) was first performed in the 1940s and is created by division of the subclavian artery (left or right) and anastomosis to the ipsilateral pulmonary artery (PA). The main advantage is that the shunt grows with the patient and the main disadvantage is loss of pulses in the ipsilateral upper limb and resulting decreased growth and strength. The classical BT shunt is seldom used these days.

The Modified BT Shunt (MBTS) is now performed and consists of a Gore-Tex (PTFE) graft (3.5-5.0 mm diameter) interposed between the innominate or subclavian artery and the ipsilateral pulmonary artery (PA). This can be performed on the left or right side, but is routinely on the right side at PMH. It is usually a non-bypass procedure and is performed through a lateral thoracotomy.



Postoperatively, pulmonary and systemic vessels are fed from the same common source. The balance of systemic and pulmonary flows is of utmost importance, and is the principal determinant of systemic oxygen saturation (SaO_2). Ideally the post-op SaO_2 is 75-80%, and this gives a $\text{Qp}:\text{Qs}$ ratio 1:1-1.8. Refer to [Cardiac: Neonatal Circulation Changes/Unbalanced Circulation](#).

Routine Post-Op Management Following Neonatal BT Shunt Insertion

For more general post-op cardiac management, refer to [Cardiac: Routine Post-Operative Care](#) guideline.

Currently, neonates undergoing BT shunt procedures are cared for on PCC post operatively, usually until extubated and stable. The following care remains relevant once the neonate returns to NICU.

Respiratory

- Will require ventilation at least overnight, however usually weaned to extubation on day 2 if otherwise stable (PCC).
- Keep SaO_2 70-85% (ideal 75-80%).
- Gradually wean oxygen therapy down to air if $\text{SaO}_2 > 80\%$ (to avoid over-shunting).
- 6 hourly blood gases if stable

CVS

- Check for shunt murmur when arrives back from PCC and again anytime when there are concerns over the patient, particularly if they have low SaO_2 .
- Aim for BP in normal range (Refer to [Monitoring and Observation Frequency](#) guideline) unless otherwise ordered.
- If inotropic support required, 1st line dopamine, 2nd line noradrenaline if low diastolic pressure/over-shunting or adrenaline to support cardiac performance.

Cardiac: Blalock-Taussig Shunt (BT Shunt) Management Following Insertion

- An echocardiogram should be performed and documented in the notes by the cardiologist if there is a concern over shunt blockage or pulmonary over-circulation.

Causes of Hypotension/Low Cardiac Output

- Hypovolaemia/bleeding.
- 'Over-shunting' causing unbalanced circulation (particularly low diastolic pressure).
- Shunt blockage.
- Sepsis

Fluids/Nutrition

- Fluids should be restricted to 60-80 mL/kg/day on day 1 (PCC). Then can then be increased by 10-20 mL/kg/day depending upon the clinical status of the patient.
- Beware of over-restriction causing intravascular depletion as this predisposes to shunt thrombosis.
- Feeds can be considered the day after surgery if everything has been stable. Grade up feeds as per normal protocol.
- If there are concerns of 'over-shunting' and a poor systemic output, feeds should be withheld as there is a risk of NEC.

Anticoagulation

- Anticoagulation with heparin to prevent BT shunt thrombus is controversial. It is not routinely used in every case at PCH. The decision to use heparin is on a case-by-case basis and is discussed with the cardiac surgeon on returning from theatre. The heparin infusion is usually commenced at a rate of 10 units/kg/hr (can use up to 20 units/kg/hr). Changes in APTT are not usually targeted.
- Aspirin 3-5 mg/kg once a day is commenced once on feeds.

Haemoglobin

- Should be kept around 120 g/L. Avoid too high levels as this may predispose to shunt thrombosis. Avoid a low Hb as patients with cyanotic heart disease require slightly higher Hb levels than normal to optimise oxygen carrying capacity.

Post-Op BT Shunt Complications – Refer to [Cardiac: Complications Management Following Surgery](#)

- **Haemorrhage** from anastomosis may lead to haemothorax.
- **Poor shunt flow/blockage** with thrombus or shunt too small (unusual – refer to [Low O₂ Saturations < 70%](#) section below).
- **'Over-shunting'** - additional blood supply from PDA (which is yet to close) or MAPCAs, or shunt too large.
- **Shunt infection** (rare) results in features of sepsis with elevated CRP and WCC.
- Discuss with cardiac surgeon **before** commencing antibiotics if this is felt to be a possibility.
- **Seroma** - Gore-Tex can 'sweat'. Seen on CXR or echo.
- **Chylothorax** due to thoracic duct damage.
- **Vocal cord palsy** due to recurrent laryngeal nerve.
- **Diaphragmatic palsy** due to phrenic nerve damage.
- **Shunt narrowing** may occur at the site of anastomosis and may sometimes need to be treated with balloon angioplasty.

Low O₂ Saturations (< 70%)

Usually due to inadequate pulmonary blood flow.

Causes

- Shunt blockage with thrombus.
- Inadequate shunt flow due to systemic hypotension.
- Small feeding artery/ pulmonary arteries (PAs)/branch PAs.
- Shunt too small (unusual as the shunt is usually larger than the feeding artery to allow for growth of the baby and so the feeding artery is usually the limiting factor).
- Inadequate ventilation e.g. Pneumothorax, atelectasis, pneumonia, haemothorax, chylothorax, dislodged ETT.

Signs

A shunt blockage may present acutely or sub-acutely and can be complete or partial. If there is an acute complete blockage it is an emergency and will present with:

- Sudden severe desaturation.
- Normal (or low) blood pressure.
- Normal ventilator parameters, normal air entry and normal CXR.
- **No shunt murmur heard** on auscultation.

Management

- If SaO₂ < 65% and shunt malfunction suspected (i.e. no respiratory cause) call NICU consultant. Inform cardiac surgeon and cardiologist immediately.
- Review of ventilation status, blood gas and CXR should have been performed.
- Ensure haemodynamic status is adequate. May require volume expansion with normal saline. Give 5 mL/kg aliquots and titrate with response. It is essential to give smaller aliquots of fluid so as not to give too much fluid which could tip the delicate balance of a cardiac patient. **If requiring > 20-30 mL/kg of fluid, reconsider your diagnosis.**
- May require vasopressors to augment systemic pressure. Noradrenaline works well by intense peripheral vasoconstriction to increase the systemic driving pressure. However dopamine and/or adrenaline may be required to support the cardiac function.
- Ensure coagulation profiles are normal and blood available if trip back to theatre seems likely.

High O₂ Saturations (> 85% in Air)

Usually due to high shunt flow, causing excessive blood flow to the lungs and an unbalanced circulation.

Causes

- Unnecessarily high FiO₂.
- PDA still open and so additional shunting of blood to the lungs*.
- MAPCAs as a source of additional blood flow to the lungs.
- Shunt too large.

*The ductus arteriosus (DA) is not routinely ligated at PMH as most procedures are right sided and access to the DA is not possible. Therefore the DA is patent post-op and causes additional blood flow to the lungs. PGE1 will have been ceased in theatre, and the PDA

Cardiac: Blalock-Taussig Shunt (BT Shunt) Management Following Insertion

usually closes over hours/days. If the high O₂ saturations are due to this reason, then they should improve as the ductus closes.

Other Signs

- **Rising lactate and metabolic acidosis in a 'pink' patient** (reduced systemic flow)
- Pulmonary plethora/oedema (may be unilateral).
- Cardiac failure.
- Systemic diastolic pressure low due to 'run off' to the lungs.
- May cause pulmonary haemorrhage.

Treatment

- Ensure ventilating in FiO₂ 0.21.
- Inform NICU consultant.
- If SaO₂ newly consistently > 90% inform cardiologist (will require review and echo)
- If not mechanically ventilated, consider ventilation and use higher PEEP/MAP.
- In more refractory cases manipulation of pH (permissive hypercapnia) may assist in increasing the pulmonary vascular resistance (Refer to [Cardiac: Neonatal Circulation Changes/Unbalanced Circulation](#)).
- Often fluid restriction and diuretics are beneficial.
- Sometimes volume loaded ventricles needs supporting with inotropes e.g. [Dopamine](#).
- If poor systemic output, withhold feeds as at high risk of NEC.
- Occasionally urgent surgical revision of a large shunt is required (smaller diameter shunt inserted).

Prior to Discharge

Whenever a child with a BT shunt develops an illness where they may become dehydrated e.g. Gastroenteritis/vomiting or has a decreased oral intake or is sweating excessively, they are at risk of shunt thrombosis. They will need increased oral fluids e.g. Oral rehydration solution (ORS) if tolerated or early IV fluids. Parents should be warned of this prior to discharge and told to seek medical help early.

Also, if their child ever becomes blue then they need to take them to hospital immediately as it may indicate shunt blockage.


Related CAHS internal policies, procedures and guidelines

Neonatology Guideline

- [Cardiac: General Complications Management Following Surgery](#)
- [Cardiac: Neonatal Circulation Changes/Unbalanced Circulation](#)
- [Cardiac: Routine Post-Operative Care](#)
- [Monitoring and Observation Frequency](#)
- [Post-Operative: Analgesia](#)

References
1. Artman, M., Mahony, L., & Teitel, D. F. (2011). <i>Neonatal Cardiology</i> (Second ed.). USA.
2. Duncan, A., & Croston, E. (2008). <i>Guidelines for the Intensive Care Management of Infants and Children after Congenital Heart Surgery</i> . Paediatric Intensive Care Unit, Princess Margaret Hospital for children, Perth.
3. Gedicke, M. et al. <i>Risk factors for acute shunt blockage in children after modified Blalock Taussig shunt operations</i> . Heart vessels 2010; 25: 405-409
4. Horrox, F. (2002). <i>Manual of Neonatal and Paediatric Heart Disease</i> (First ed.). Gateshead, Tyne and Wear, UK: Whurr Publishers.
5. Koenig, P., Hijazi, Z. M., & Zimmerman, F. (2004). <i>Essential Pediatric Cardiology</i> (First ed.). USA: McGraw-Hill Companies, Inc.
6. Nichols, D. G., Ungerleider, R. M., Spevak, P. J., Greeley, W. J., Cameron, D. E., Lappe, D. G., & Wetzel, R. C. (2006). <i>Critical Heart Disease in Infants and Children</i> (Second ed.). USA: Mosby.
7. Swain, K. S. et al. <i>Neonatal Blalock-Taussig Shunt: Technical aspects and postoperative management</i> . Asian Cardiovasc and thorac ann: 2008; 16: 7-10

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Neonatology / Cardiology		
Reviewer / Team:	Neonatal Coordinating Group		
Date First Issued:	May 2013	Last Reviewed:	26 th May 2020
Amendment Dates:		Next Review Date:	26 th May 2023
Approved by:	Neonatal Coordinating Group	Date:	23 rd June 2020
Endorsed by:	Neonatal Coordinating Group		
Standards Applicable:	NSQHS Standards: 		

Printed or personally saved electronic copies of this document are considered uncontrolled



Healthy kids, healthy communities

Compassion
Excellence
Collaboration
Accountability
Equity
Respect

Neonatology | Community Health | Mental Health | Perth Children's Hospital