

Oral Glucose Tolerance Test

Background

This test is done to diagnose possible diabetes mellitus. It measures the change in blood glucose after a known amount of glucose has been taken by mouth.

Key points

Please ensure your child:

- Has a normal diet and normal level of physical activity for three days before the test.
- Fasts from midnight (or such time as advised by their doctor) before the test.
- It is fine for them to drink water.
- Remains fasting until the end of the test.

If your child is unwell (fever, vomiting or diarrhoea), please phone the Day Treatment Unit on 6456 3772 to cancel and rebook.

The test will be arranged for first thing in the morning.

Arrival

Please go to the Day Treatment Unit. It is on the first floor at Clinic E. Take the Yellow or Pink lifts.

The Oral Glucose Tolerance Test

Once admitted a local anaesthetic cream will be applied to your child's hand and/or inner elbow. This cream takes about an hour to work. An intravenous line can then be put in without causing any pain.

A needle is used to insert the intravenous cannula. Once in the vein the needle is taken out. A plastic tube is left lying in the vein. This plastic tube should not cause any pain. Once in, it stays in for the whole test.

To start a blood test/ tests are taken. Your child is then given a flavoured sweet glucose drink (300ml) to be swallowed within five minutes. Your child will need to stay seated throughout the test.

Possible side effects are nausea and vomiting.



Blood samples are taken at one hour and two hours after the glucose drink. The blood samples are taken using the plastic tube. The blood samples are then sent for testing. The test will take two hours.

*Your doctor may order a number of other blood tests to be taken at the start of the test to check such things as liver function, fats in the blood and insulin levels.

After all the blood samples are taken, your child will be given lunch, the intravenous line will be removed and you can go home if your child is otherwise well.

Results

The results will be sent to your child's doctor in two to three weeks. You will receive these results at your next Endocrine Clinic or private endocrine appointment, or your child's doctor may phone you.

Disclaimer: The test requested by your child's doctor requires your child to consume a glucose drink which contains food additives approved for use in Australia. These additives are also used in many sweet food items such as muesli bars and fruit juices. Allergic reactions to these additives are very rare however if you think your child may be allergic to any of the following you should inform the collection staff: 202 (Sorbate): 211 (Benzoate): 330 (Citric acid): 143 (Fast Green).

This document can be made available in alternative formats on request for a person with a disability.

Child and Adolescent Health Service
15 Hospital Avenue, Nedlands, WA, 6009
Telephone: (08) 6456 2222
Produced by Endocrinology and Diabetes
Ref: 846 © CAHS 2020



Government of **Western Australia**
Child and Adolescent Health Service

Disclaimer: This publication is for general education and information purposes. Contact a qualified healthcare professional for any medical advice needed.
© State of Western Australia, Child and Adolescent Health Service