



GUIDELINE

Factors impacting on child health and development

Scope (Staff):	Community health
Scope (Area):	CAHS-CH, WACHS

Child Safe Organisation Statement of Commitment

CAHS commits to being a child safe organisation by applying the National Principles for Child Safe Organisations. This is a commitment to a strong culture supported by robust policies and procedures to reduce the likelihood of harm to children and young people.

This document should be read in conjunction with this [disclaimer](#)

Contents

Aim.....	1
Risk.....	2
Background.....	2
Principles.....	3
Key points.....	3
Clinical Practice Implications.....	4
Appendix A: Alcohol and other drugs.....	9
Appendix B: Culture and health.....	15
Appendix C: Disability and Development.....	22
Appendix D: Disadvantage.....	25
Appendix E: Family domestic violence or conflict.....	32
Appendix F: Family Mental Health.....	37
Appendix G: Homelessness, transience and/or overcrowding and remoteness.....	43
Appendix H: Trauma.....	48

Aim

To provide community health staff with information to support children and their families experiencing adverse factors and circumstances that may impact on children’s health and developmental outcomes.

Risk

Children and their families may not receive the service they need and will be at increased risk for poorer health and developmental outcomes.

Background

Child health services aim to improve the health, development and wellbeing of children and families through a model of progressive universalism. Services are offered to reflect universal health provision and include services and care planning that is proportionate to client need.

It is recognised that the compounding effect of a number of concerns may increase the level of risk for children and increase a family's vulnerability to negative outcomes. Consequently, these families would benefit from the additional services offered as Partnership contacts and where relevant, targeted referrals. A greater investment in working with families who need a higher level of support will improve long term health outcomes for children and families.

The environments in which a child develops, and its characteristics can shape different aspects of a child's development by increasing their risk of experiencing poor developmental outcomes or conversely helping them build protection from the impact of adversity.¹

There is growing evidence on the short and long term risks to health and wellbeing as a result of adverse life experiences in children, particularly when the 'adversities are prolonged, cumulative, or occurring during sensitive periods in early neurobiological development'.²

Children and young people's probability of experiencing adverse health outcomes is the interaction between risk factors (circumstances, events that increase the likelihood of poor outcomes) and protective factors (factors that moderate risk and promote healthy development/wellbeing).^{3, 4}

Protective factors contribute to providing a physical and psychosocial environment that enable children and families to feel strong and resilient and in which a child might achieve optimal growth, development and wellbeing.⁴

Risk factors that may increase a child experiencing adverse health outcomes and the need for additional services include:^{5, 6}

- trauma
- family and domestic violence or conflict
- family alcohol or drug misuse
- homelessness, transience and/or overcrowding and remoteness
- severe or untreated household member mental health issue
- child with disability or significant developmental delay
- exposure to criminality/criminal behaviour
- social isolation and exclusion
- disadvantage
- racial discrimination
- sexual abuse

It is important to note that not all families with complex concerns or risk factors will require additional services, as the presence of protective factors may reduce adversity and increase resilience. Protective factors can mitigate the impact of risk factors and include:^{1, 3, 6-9}

- positive parent/care giver- child relationship and kinship care
- nurturing and secure attachment
- family stability and support
- knowledge, attitudes and beliefs
- connection to community/family
- connection to culture
- mother's education level
- high self-esteem and resilience
- social support (mental health)
- social and emotional competence of child
- connection to land and country

Risk factors can accumulate for children and young people across the life span and at key transition points (e.g. starting school, transition into high school). Enhancement of protective factors in care planning and service delivery can support positive outcomes.⁶

Principles

- Services are offered through a model of progressive universalism and include services and care planning that is that is proportionate to client need.
- Consideration of protective factors supports a strengths-based family centred approach which enhances engagement with the family.
- The nurse and clients will work together for the shared understanding of family concerns and resilience, and the establishment of goals to facilitate change for modifiable concerns.
- Providing services according to need and circumstances of client and the understanding that not all families with complex concerns will require additional services, as the presence of protective factors may reduce adversity and increase resilience.
- For Aboriginal children and their family protective factors such as connection to community and culture, kinship and a sense of belonging can positively influence health outcomes.⁷⁻⁹

Key points

- The child is the primary client and is the centre of care
- Family-centred and strengths-based approaches are used, for a shared understanding of concerns and care planning that is proportionate to client needs

- The wellbeing of families and children might be adversely impacted by individual, parental or family circumstances at different time points; creating risk of poor health or developmental outcomes
- Being exposed to risk doesn't always lead to poor outcomes, protective factors can lessen children's risk of adverse health and developmental outcomes
- All nurses will refer to the [Nursing and Midwifery Board AHPRA Decision-making framework](#) in relation to scope of practice and delegation of care to ensure that decision-making is consistent, safe, person-centred and evidence-based.

Clinical Practice Implications

Community nurses play an important role in supporting clients and their families who are experiencing factors that may impact on health and developmental outcomes.

Nurses are encouraged to adopt a child and family centred care approach to care. Both the child and their family should be active participants in their health care journey.

The below table provides an overview of clinical practice implications across multiple child health and development factors.

- Alcohol and other drugs
- Culture and health
- Disability and development
- Disadvantage
- Family domestic violence or conflict
- Family Mental Health
- Homelessness, transience and/or overcrowding and remoteness
- Trauma

Information on the above factors impacting child health and development can be accessed [here](#).

Clinical practice implications across the above listed factors are taken into consideration in the table below.

Area	Information
Relevant screening/assessment	Where appropriate community health nurses should: <ul style="list-style-type: none"> • update the electronic health record with appropriate information/flag to advise other staff of the specific circumstances of client and their families • ensure referrals to appropriate services are made immediately

	<ul style="list-style-type: none"> • follow up with clients and their families to ensure any referrals made have been actioned <p>Nurses should refer to the Clinical Nursing Policy Manual and follow processes described in the relevant policy for guidance, screening and assessment information including:</p> <ul style="list-style-type: none"> • Ages and Stages Questionnaire™ guideline for guidance on identifying developmental delays, determining follow-up and referral actions required, and relevant resources. • Family and domestic violence protocol for guidance on identifying FDV and determining actions required, including screening and assessment • Perinatal and Infant Mental Health guideline for information on the administration of the Edinburgh Postnatal Depression Scale (EPDS) • Sexual Assault Response procedure including information on Mandatory Reporting of Child Sexual Abuse • Guidelines for Protecting Children 2020 for information to appropriately address child abuse concerns identified through the provision of health services. • Clients of Concern management protocol for information relating to the identification and support of families with complex needs <p>Forms</p> <ul style="list-style-type: none"> • Child Wellbeing Guide- tool to assist health professionals to identify neglect and take appropriate action
<p>Relevant care planning considerations</p>	<p>Parents and carers are key partners in assessment and planning. They are the experts about their child’s functioning, and about family history, concerns, the surrounding environment, and current supports.</p> <p>Services must be provided at a level proportionate to client need.</p> <p>Nurses should consider the client’s circumstances during care planning and when referral to services are made. This includes;</p>

	<ul style="list-style-type: none"> • the complexity of the client’s circumstance (noting that it can be a combination of factors impacting the current circumstance e.g. homelessness, socioeconomic disadvantage, FDV) • that complex client and family presentations often involve the concurrent and cumulative effect of social determinants, which may require referrals to outside agencies such as housing and social services • being aware of the Principals of Trauma Informed Care and Practice when working with children and their families • offering additional contacts to meet individual needs where clinical judgement warrants • an awareness that long term care planning and support may be appropriate for some clients. • acknowledging client’s barriers to accessing services in care planning considerations <p>Consideration of the above enhances the ability and capacity of the client and their family to access necessary services within a timely period, therefore, optimising their health and wellbeing outcomes.</p> <p>Nurses can support families to take a family centred approach in their children’s health care. The consideration and enhancement of protective factors when assessing, consulting or care planning can support positive outcomes for the child and family. See individual factor papers for specific protective factors.</p>
Training requirements	Nurses are required to complete training specific to their role and local area need as per the CAHS-CH Nurses working in Child Health Nursing Practice Framework or the WACHS Nursing and Midwifery Learning Framework and Guidelines and associated individual global learning plans.
Awareness of local service availability	Community Health Nurses can assist clients and their families by being knowledgeable of local services (e.g. services that provide emotional, financial and practical assistance) that may provide support.

	Aboriginal staff within your service may be of assistance with identifying culturally appropriate local services.
--	---

References	
1.	Legge E. Risk and protective factors in early childhood: An ecological perspective [CoLab Evidence Report]. 2018
2.	Masten AS, Barnes AJ. Resilience in Children: Developmental Perspectives. <i>Children</i> 2018;5:98
3.	Commissioner for Children and Young People. Improving the odds for WA's vulnerable children and young people. In: Commissioner for Children and Young People WA, ed. Perth; 2019
4.	Child Welfare Information Gateway. Issue Briefs: Protective Factors Approaches in Child Welfare. 2020
5.	Rickwood D, Thomas K. Mental wellbeing interventions: an Evidence Check rapid review brokered by the Sax Institute for VicHealth. Victoria; 2019
6.	Department of Health. National Action Plan for the Health of Children and Young People 2020-2030. Commonwealth of Australia; 2019
7.	Hunter S-A, Skouteris H, Morris H. A Conceptual Model of Protective Factors Within Aboriginal and Torres Strait Islander Culture That Build Strength. <i>Journal of Cross-Cultural Psychology</i> 2021;52:726-751
8.	Australian Institute of Health and Welfare. Aboriginal and Torres Strait Islanders Health Performance Framework- Measures. 2020. Available at: https://www.indigenoushpf.gov.au/measures2021
9.	Australian Institute of Health and Welfare. Aboriginal and Torres Strait Islander Health Performance Framework 2020 key health indicators—Western Australia. Canberra: AIHW; 2020


Related internal policies, procedures and guidelines
The following documents can be accessed in the Community Health Manual: HealthPoint link or Internet link or for WACHS staff in the WACHS Policy link
Children in Care- conducting an assessment
Children in Care- managing referrals for assessment
Clients of concern management
Family and domestic violence
Guidelines for Protecting Children 2020
Partnership- child health service

Perinatal and infant mental health
Sexual assault response
Universal Contact suite
Universal Plus- Child Health

Related external legislation, policies, and guidelines (if required)
External Legislation, Standards and Policy (list and hyperlink)
Mandatory reporting of child sexual abuse

Useful internal resources (including related forms) (if required)
Child Wellbeing Guide 0-18 Years (CHS470)

This document can be made available in alternative formats on request.

Document Owner:	Nurse Co-Director, Community Health		
Reviewer / Team:	Clinical Nursing Policy Team		
Date First Issued:	19 July 2022	Last Reviewed:	
Amendment Dates:		Next Review Date:	19 July 2022
Approved by:	Community Health Clinical Nursing Policy Governance Group	Date:	24 June 2022
Endorsed by:	Executive Lead Community Health	Date:	14 July 2022
Standards Applicable:	NSQHS Standards:  1.15, 2.6, 2.7, 2.9, 2.10, 5.7, 5.9, 5.10, 5.12, 5.13, 5.14, 6.3 Child Safe Standards: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10		

Printed or personally saved electronic copies of this document are considered uncontrolled



Healthy kids, healthy communities

Compassion
Excellence
Collaboration
Accountability
Equity
Respect

Neonatology | Community Health | Mental Health | Perth Children's Hospital

Appendix A: Alcohol and other drugs

Key Points

- Parental substance misuse occurs on a continuum and can have negative physical, developmental, psychosocial and emotional impacts on the child across the life course
- Timing, length and number of exposures can affect the severity of impact to the child
- The most commonly misused substance in Australia is alcohol
- Parental substance misuse commonly occurs in the presence of multiple complex issues, such as mental illness, family and domestic violence, homelessness and poverty. Children are more likely to be negatively impacted where mental health issues and/or family and domestic violence are present
- The impact of parental substance misuse can vary between children and protective factors may mitigate negative impacts


Definition

Alcohol and other drugs are psychoactive substances which act on the central nervous system and alter the way a person thinks, acts and behaves ¹. They can include illegal, prescription or over-the-counter substances, for example:

- Depressants e.g. alcohol, valium, xanax, GHB, kava
- Stimulants e.g. speed, cocaine, ice
- Opioids e.g. heroin, fentanyl, buprenorphine, morphine, oxycodone, codeine
- Psychadelics e.g. LSD, magic mushrooms, DMT
- Cannabinoids e.g. synthetic/medicinal cannabis, marijuana, butane hash oil
- Dissociatives e.g. nitrous oxide, ketamine
- Empathogens e.g. Ecstasy, PMA, bath salts

Substance misuse or abuse can be defined as ‘the harmful or hazardous use of psychoactive substances, including alcohol and illicit drugs’ ². It is important to recognise that substance use exists on a continuum and can be problematic, regardless of whether or not a person is dependent on the substance ¹.

Spectrum of substance use ³



Signs that a person's substance use is problematic may include ³:

- Regularly using more of the substance than intended
- Requiring increasing quantities of the substance to achieve the desired effect
- Spending too much time acquiring, using or recovering from the substance
- Difficulties meeting work, school or family responsibilities
- Experiencing physical symptoms when not using the drug which are only relieved by taking the drug
- Relationship problems caused by use of the drug
- Failed attempts at cutting down or stopping use of the drug
- Cravings or strong desire to use the substance

Prevalence

Around 1 in 20 Australians has an addiction or substance use problem, with alcohol being the most commonly misused substance ⁴. In 2019, the most commonly used illicit substance was cannabis (11.7%), followed by cocaine (4.2%), ecstasy (3%) and opioids (2.7%) ⁵ Approximately 35% of Australians with a substance use disorder also experience a mental health disorder ⁵.

Alcohol is consumed by 77% of Australians ⁶. 1 in 3 alcohol drinkers exceed national guidelines by drinking more than 4 standard drinks in one sitting, a level attributed to increased risk of violence, assault, motor vehicle crashes and other injury ⁶. Alcohol was involved in 29% of all family violence incidents in 2019 ⁶. Children with parents or guardians who experience alcohol dependence are more likely to be brought to the attention of child protective services ^{6,7}. In 2006-2007, it is estimated that 20,000 Australian children were victims of alcohol related child-abuse ⁶.

Health and developmental impacts/outcomes for child

All areas of a child’s life can be negatively impacted by problematic parental substance use across the life course (see Table 1). ^{8,9} It is important to note that all children are impacted differently by parental drug use. ^{8,9}

Parental substance misuse often co-occurs with many other complex issues ¹⁰ The presence of additional issues increases risk of impact to the child, particularly parental mental illness and exposure to domestic violence ^{8,9,11} Children of parents who misuse substances are at increased risk of maltreatment, neglect and abuse. ^{7,8,9,11}

The risk of impact to the child is cumulative in accordance with the number of factors and length of exposure. ⁸ The presence of problems at key development stages during early life is thought to be particularly influential. ⁸

Additional risk factors include:

- Parental mental illness
- Domestic violence/abuse
- Poverty and socioeconomic disadvantage
- Unemployment
- Homelessness/housing instability
- Social exclusion and discrimination
- Family disruption, separation and substitute care
- Criminal activity
- Absence of stable adult figure

Ways in which parental substance misuse can affect children across the life course include ¹²:

- Direct physiological effects (eg: foetal exposure to substance)
- Direct harm to a child by the intoxicated parent
- Diversion of parental attention to substance using activities
- Parental modelling of substance misuse behaviours to the child

Table 1: Health and developmental impacts of problematic parental alcohol and drug-use to children across the life-course ^{9, 11, 13-15}

Age of child	Impacts/outcomes
Unborn child	<ul style="list-style-type: none"> • Spontaneous abortion • Miscarriage • Placental abruption • Prematurity • IUGR • SGA • Low birth-weight

	<ul style="list-style-type: none"> • Foetal Alcohol Spectrum Disorder • Neurological damage • Neonatal Abstinence Syndrome • Irritable temperament • Poor parental attachment
Babies/toddlers/pre-schoolers	<ul style="list-style-type: none"> • Poor growth • Nutritional deficiencies • Increased risk of SIDS • Poor parental attachment • Increased risk of accidents • Cognitive deficit
School-aged/pre-adolescents	<ul style="list-style-type: none"> • Mood/anxiety disorders • Anxiety and depression • Aggression & withdrawal • Inattentiveness • Hyperactivity • Behavioural/attentional disorders • Cognitive deficit • Impaired academic achievement • Impaired social functioning
Adolescents	<ul style="list-style-type: none"> • Substance use problems • Mood/anxiety disorders • Anxiety and depression • Aggression & withdrawal • Impaired academic achievement • Impaired social functioning

Protective Factors

Whilst children who are exposed to parental substance misuse have a greater risk of negative outcomes, protective factors can reduce impact (see Table 2) ⁸

Individual factors	<ul style="list-style-type: none"> • Internal locus of control • Self-monitoring/coping skills/self-control • Effective emotional expression • Hobby/creative outlet • Future planning • Good understanding of parental misuse behaviour • Intellectual capacity • Abstinence from alcohol and drugs
Family factors	<ul style="list-style-type: none"> • Supportive relationship with a stable (non-substance misusing) adult • Demonstrations of affection from extended family • Parental self-efficacy/self-esteem • Consistency and stability in every-day/family life

	<ul style="list-style-type: none"> • Constructive coping styles and deliberate actions by parents to minimise adversity for children • Strong family norms and morality • Adequate finances and good employment opportunities • Positive care style of parents • Parental modelling of behaviours expected from child • Absence of domestic violence/abuse
Parental factors	<ul style="list-style-type: none"> • Parental problems are of reduced severity and shorter duration • One parent does not have problems • Parent is receiving treatment • Drug activity/paraphernalia is kept hidden/drug use occurs away from the home
Community /environmental factors	<ul style="list-style-type: none"> • Cultural connectedness, values and identity • Support from community adult role model eg: teacher, neighbour • Strong friendships/peer relationships • Positive school experiences/consistent attendance at school • Support from key community services eg: healthcare

References	
1.	Alcohol and Drug Foundation. Drug Facts 2021 [Available from: https://adf.org.au/drug-facts/].
2.	World Health Organization. Substance Abuse 2021 [Available from: https://www.afro.who.int/health-topics/substance-abuse#:~:text=Substance%20abuse%20refers%20to%20the%20harmful%20or%20hazardous,the%20negative%20health%20consequences%20experienced%20by%20its%20members.]
3.	NSW Government. Handbook for nurses and midwives: Responding effectively to people who use alcohol and other drugs. In: Health N, editor. St Leonards NSW 2021.
4.	HealthDirect. Substance Abuse 2020. Available from: https://www.healthdirect.gov.au/substance-abuse .
5.	Australian Institute of Health and Welfare. National Drug Strategy Household Survey 2019.
6.	Australian Department of Health. National Alcohol Strategy. 2019.
7.	Australian Institute of Health and Welfare. National framework for protecting Australia's children indicators 2021.
8.	Velleman R, Templeton LJ. Impact of parents' substance misuse on children: an update. BJPsych Advances. 2018;22(2):108-17.
9.	Horgan J. Parental substance misuse: Addressing its impact on children. National Advisory Committee on Drugs; 2011.
10.	Dawe S, McMahon TJ. Innovations in the Assessment and Treatment of Families with Parental Substance Misuse: Implications for Child Protection. Child Abuse Review. 2018;27(4):261-5.
11.	Smith VC. Families affected by parental substance use. American Academy of Paediatrics. 2016;138(2).

Factors affecting child health and development

		<p>IDA, referral to dentist for children, treatment plan (and health promotion advice) in place for sores and scabies</p> <ul style="list-style-type: none"> • Whole family health check 	<p>increase iron levels</p> <ul style="list-style-type: none"> • Nutrition advice and/referral to food help agency • Referral to a public general dental clinic 	<ul style="list-style-type: none"> • Medical Officer/GP • Aboriginal Health Team staff (AHW & CN)
Mind and Emotions	<ul style="list-style-type: none"> • AHT 	<ul style="list-style-type: none"> • 4 & 3-year-old are engaged with the nearest Kindilinks/Child and Parent Centre • ASQ assessment • EDPS 	<ul style="list-style-type: none"> • Work with Jodie's 3-year-olds behavioural issues • Parents and Aunty provided with strategies to support 3-year-old 	<ul style="list-style-type: none"> • CDS staff • AHW • Kindilinks
Family and Kinship	<ul style="list-style-type: none"> • AHT 	<ul style="list-style-type: none"> • Jodie and father connected into family support/parenting programs • Strengthen support from Aunty and older cousins 		<ul style="list-style-type: none"> • Aunty and other parents • Family • Family support programs
Community	<ul style="list-style-type: none"> • AHT 	<ul style="list-style-type: none"> • Kindilinks • Linking in with other services (Public Housing, financial counselling) 		<ul style="list-style-type: none"> • Community • Kindilinks
Culture	<ul style="list-style-type: none"> • Cultural events 	<ul style="list-style-type: none"> • Encourage Jodie to attend cultural events with children 		
Country				
Spirit				

References	
1.	James M. Aboriginal and Torres Strait Islander health performance framework 2014 report. 2015
2.	Bryant J, Bolt R, Botfield J, et al. Beyond deficit: 'strengths-based approaches' in Indigenous health research. <i>Sociology of health & illness</i> 2021;43:1405-1421
3.	Healing Foundation. A Theory of Change for Healing. 2019

4. Department of Health. National Framework for the Health Services for Aboriginal and Torres Strait Islander Children and Families. Canberra: Australian Government; 2016
5. Gee G, Dudgeon P, Schultz C, et al. Aboriginal and Torres Strait Islander social and emotional wellbeing. *Working together: Aboriginal and Torres Strait Islander mental health and wellbeing principles and practice* 2014;2:55-68
6. Department of Health. National Aboriginal and Torres Strait Islander Health Plan 2021-2031. Australia; 2021
7. First 1000 Days Australia. Why First 1000 Days?
8. Fogarty W, Lovell M, Langenberg J, et al. Deficit discourse and strengths-based approaches. *Changing the Narrative of Aboriginal and Torres Strait Islander Health and Wellbeing Melbourne: The Lowitja Institute* 2018
9. Australian Institute of Health and Welfare. Aboriginal and Torres Strait Islander Health Performance Framework 2020 summary report. Canberra: AIHW; 2020
10. Houston S. A national Aboriginal health strategy. *Aboriginal and Islander Health Worker Journal* 1989;13:7-8
11. Western Australia Office of Aboriginal Health. Aboriginal cultural security: a background paper. Perth; 2003
12. Australian Indigenous Health Infonet. Cultural practices.
13. Dudgeon P, Milroy H, Walker R. Working together: Aboriginal and Torres Strait Islander mental health and wellbeing principles and practice. Telethon Kids Institute, Kulunga Aboriginal Research Development Unit.; 2014
14. Swan P, Raphael B. Ways Forward. Canberra: Australian Govt. Publishing Service; 1995
15. Government of Western Australia Department of Health. WA Aboriginal health and wellbeing framework 2015–2030. 2015
16. Australian Health Ministers' Advisory Council. Cultural Respect Framework 2016-2026 for Aboriginal and Torres Strait Islander health. Author Canberra; 2016
17. Department of the Prime Minister and Cabinet. National Strategic Framework for Aboriginal and Torres Strait Islander Peoples' Mental Health and Social and Emotional Wellbeing 2017-2023. Canberra; 2017

Appendix C: Disability and Development

Key points

- Disability is the interaction between a child's health conditions, body functions and structures, the activities they participate in, and the environmental factors that affect these.^{1,2}
- A child may be born with disabilities, or these may develop after birth. Disabilities may be caused by genetics, injury or illness.³
- Children with disability may have special needs, and may require early intervention and increased support.³
- Children with intellectual disability or mental and behavioural problems have a greater risk of experiencing maltreatment than children without disability.⁴
- The health and development of children with disability can be improved by timely surveillance, thorough assessment, early referral for services, early intervention, inclusive schooling, and support for families/carers.^{5,6}

Definitions

Disability is defined by ABS as “any limitation, restriction or impairment which restricts everyday activities and has lasted, or is likely to last, for at least six months”.⁷ It refers to any condition that restricts a person’s mental, sensory, intellectual or mobility functions and may be caused by accident, genetics or disease.⁵ The impairment can be temporary or permanent, total or partial, visible or invisible, and can be there from birth or occur during a person’s lifetime.

Developmental delay describes a lag in the acquisition of a skill or milestone otherwise expected of a child at a particular age.⁸ Developmental delays are measured using validated developmental assessments and may be mild, moderate or severe. While a developmental delay may not be permanent, it can provide a basis for identifying children who may experience a disability.⁵

Developmental disability describes the profile of children with complex and pervasive developmental difficulties that are likely to impact on a child’s ability to participate optimally in functional activities across their lifecourse.⁸

Neurodevelopmental disorders are disorders of early brain development. They include autism spectrum disorder (ASD), intellectual disability, motor disability (e.g. cerebral palsy), seizures, learning disabilities (e.g. dyslexia), and attention deficit hyperactivity disorder (ADHD). Children with neurodevelopmental disorders can experience a wide range of symptoms, including reduced emotional regulation, poor movement (motor) control, problems with language development and social integration, and impacted learning ability.⁹

Limitation means a person has difficulty, needs assistance from another person, or uses an aid or other equipment to perform one or more of the core activities (communication, mobility and self-care). The severity of limitations can be mild, moderate, severe or profound.⁴

Schooling restriction means a child needs special assistance or equipment to participate in a mainstream class or attend a special school or special classes.⁴

Prevalence

In WA in 2018, 7.5 % (43,600) of children and young people aged 0 - 17 years had a reported disability.⁷ Around 11.5% (22,400) of children in WA aged 6 - 11 years have a disability.⁷ Disabilities in childhood are varied, and include cerebral palsy, intellectual disability, spina bifida, acquired brain injury, visual or hearing impairment, autism spectrum disorder, and rare genetic conditions such as tuberous sclerosis.⁹ Boys are almost twice as likely as girls to have a disability.⁴

Across Australia, 70 % of 0 - 5-year-old children with disability have a speech or sensory disability (including loss of sight or hearing). The most common types of disability in the 6 - 11 year-old age group are intellectual (67.8%) and psychosocial (39.3%) disability.⁷

Just over half (52%) of children with a disability have a profound or severe core-activity limitation and require assistance with one or more core activities of daily.¹ The 2016 Census estimated that Indigenous children aged 0 – 14 years were 1.7 times as likely as non-Indigenous children to have a severe core-activity limitation, but considered this to be a significant underestimation.⁴

In 2017, 18.8% of all Australian primary and secondary students received an adjustment at school to address disability.⁴ Almost all (97%) children aged 5–14 years with a disability were attending school; 89% were in mainstream schools and 9% were in schools specially designed for students with disability.³

The increase in prevalence rates for autism and ADHD has been attributed to improving diagnostic methods and increased awareness.¹⁰

Health impacts and outcomes

The impairments related to a disability may interact with various barriers to hinder a child's full, effective and equal participation in society.¹¹ To reach their full potential, all children need good health care, nutrition and safety, responsive care giving, early learning opportunities, inclusive schooling, and opportunities to take part meaningfully in home and community activities. Children with disability have all the same needs but may require extra support to help them have these needs met.

Development proceeds through a series of milestones. Typically, simple skills are mastered before more complex skills can be learned. Developmental delays or disabilities in one area can impact on the child's ability to consolidate skills and progress through to the next developmental stage. Chronic health conditions can also have long-term effects on a child's development and behaviour.¹² The broad range of individual differences between children often makes it difficult to distinguish between typical variations in development, maturational delays, transient disorders, and persistent impairments.⁸

Children with disability are at risk of the same childhood illnesses as other children. They may have specialised health-care needs related to their disability, and other secondary conditions. For example, children who are wheelchair users are vulnerable to pressure ulcers.⁵

Children with disability can be disproportionately exposed to risk factors such as poverty, stigma and discrimination, poor caregiver interaction, violence, abuse and neglect, and limited access to programmes and services. All can have a significant effect on their wellbeing and development.⁵

Families of children with disabilities experience more stress, greater financial strain and poorer wellbeing than families with typically developing children.¹³ This is particularly due to the time and emotional commitments associated with raising a child with high support needs. These parents have an increased risk of developing mental health problems such as depression and anxiety, and significant stress on familial and social relationships.¹³

Mothers have described emotional support as possibly the most important influential coping factor.¹ Support is most critical at the time of diagnosis and during medical intervention for their child.¹⁴

Protective factors

Effective interventions can alter the course of a child's development by positively changing the balance between protective and risk factors within a child's environment.⁸

Protective factors that may reduce the incidence and severity of impact of disability and developmental delay in children include:

- a sense of belonging to home, family and community, and a strong cultural identity
- pro-social peer group
- positive parental expectations and home learning environment
- positive opportunities at major life transitions
- access to child and adult focused services, including general and mental health, maternal and child health, early intervention, disability, drug and alcohol, family support, family preservation, parenting education and recreational facilities
- accessible and affordable childcare and high-quality preschool programs
- inclusive community neighbourhoods/settings
- the service system's understanding of neglect and abuse.¹²

Early intervention

Early intervention (EI) is specialised support for children with disability, autism or other additional needs including developmental delay. EI refers to therapies and supports for children and their families in the early years from birth until children start school.⁶ The World Health Organization recommends EI as the best way to support the development and wellbeing of children with disability, autism or other additional needs including developmental delay.⁵ With family involvement and timely interventions, EI helps children develop the skills they need to take part in everyday activities, and promotes a more stimulating and protective environment.⁵ Sometimes children who get EI need less or no support as they get older.⁶ There are high economic returns on early intervention, particularly for disadvantaged children, but EI must be followed up along the life course or the economic returns diminish.⁸

There is evidence that providing support and services for infants and young children with early developmental impairments and their families can alter the child's longer-term developmental trajectory and reduce the risk of secondary health and psychosocial complications. Supporting the family is a crucial component of EI programs, as the family has a key role in fostering their child's developmental potential and may experience additional stresses as they meet the special needs of their child.²

References

1. Australian Institute of Health and Welfare. Children with Disabilities in Australia.: AIHW; 2004:117
2. Royal Australasian College of Physicians - Paediatric and Child Health Division. Position Statement: Early Intervention for Children with Developmental Disabilities. 2013. Available at: https://www.racp.edu.au//docs/default-source/advocacy-library/early-intervention-for-children-with-developmental-disabilities.pdf?sfvrsn=f6a32f1a_12
3. Pregnancy Birth and Baby. What is a childhood disability? 2021. Available at: <https://www.pregnancybirthbaby.org.au/what-is-a-childhood-disability>
4. Australian Institute of Health and Welfare. Australia's Children. Canberra: AIHW; 2020
5. World Health Organization. Early childhood development and disability: A discussion paper. 2012
6. Raising Children Network. Early intervention for children with disability, autism or other additional needs. 2021. Available at: <https://raisingchildren.net.au/disability/services-support/services/early-intervention>
7. Commissioner for Children and Young People. Profile of Children and Young People in WA – 2022. Perth; 2022
8. Queensland Department of Health. Child Development in Queensland Hospital and Health Services: Act Now for kids 2morrow: 2021 to 2030. 2013
9. Murdoch Children's Research Institute. Neurodevelopment. 2021. Available at: <https://www.mcri.edu.au/research/flagships/neurodevelopment>. Accessed 21/02/2022, 2022
10. Australian Institute of Health and Wellbeing. Disability updates: children with disabilities. Canberra: AIHW; 2006
11. Australian Network on Disability. Disability statistics. 2021. Available
12. Government of Western Australia. Child development and trauma guide. In: Department for Communities, ed.; 2021
13. Carers Australia. How Does Having A Child With A Disability Affect A Family? . 2020. Available at: <https://www.carersaustralia.com.au/wp-content/uploads/2020/08/How-Does-Having-a-Child-with-a-Disability-Affect-a-Family.pdf>
14. Gilson K-M, Davis E, Johnson S, et al. Mental health care needs and preferences for mothers of children with a disability. *Child: care, health and development* 2018;44:384-391
15. Children and Young people with Disability Australia. Building strong families. 2020. Available at: https://www.cyda.org.au/images/pdf/eci_5_building_strong_families.pdf

NB - Appendix C amended on 28 September 2022

Appendix D: Disadvantage

Key Points

- Disadvantage arises through the overlapping of many factors (e.g. unemployment, low income) and the effects of the living conditions of individuals (e.g. weak social support networks, lack of opportunities, social exclusion).
- SEIFA is a widely used measure of advantage and disadvantage in a geographical area.
- Disadvantage can negatively affect children's health and developmental outcomes.

Definition

Disadvantage is a complex notion which involves many aspects of people's lives. It consists of many different dimensions, and there is no one agreed definition or way of measurement.¹ Disadvantage is not only about having a low income, but includes a lack of opportunities to participate fully in society. Several broader concepts have been used to view and measure disadvantage.¹ These include:

- poverty
- deprivation
- capabilities, and
- social exclusion.¹

In Australia, the **Socio-Economic Indexes for Areas (SEIFA)** are commonly used to measure and rank the advantage and disadvantage of Local Government Areas (LGAs). It has been created by the Australian Bureau of Statistics (ABS).^{2, 3}

Socio-economic advantage and disadvantage can be defined as people's access to material (e.g. income, housing) and social resources (e.g. support networks), and their ability to participate in society.²

SEIFA is made up of four indexes.³ These are:


- The Index of Relative Socio-Economic Disadvantage - uses information such as low income, low education and occupational status as markers of disadvantage.
- The Index of Relative Socio-Economic Advantage and Disadvantage - includes measures of advantage, as well as the information in the above index.
- The Index of Economic Resources - focuses on peoples' and households' level of access to economic resources.
- The Index of Education and Occupation – focuses on the general level of educational and occupational skills of people within an area.⁴

Each index is a summary of different information from the five yearly Census (such as income, educational level, and employment) and focuses on a different aspect of

socio-economic advantage and disadvantage. The most widely used index is the Index of Relative Socio-Economic Disadvantage.

Advantaged and disadvantaged Local Government Areas are spread throughout Australia. The most advantaged LGAs (quintile (population group) 5) tend to be located around capital cities and more coastal areas. The most disadvantaged LGAs (quintile 1) tend to be in mostly regional and rural areas.²

According to the 2016 Census, the proportion of persons in WA by index of relative socioeconomic advantage and disadvantage quintiles are as follows:²

	Quintile	Percentage %
<p>Most disadvantaged</p>  <p>Most advantaged</p>	1	13.3
	2	18.4
	3	22.9
	4	24.0
	5	21.4

SEIFA is commonly used to:

- determine which areas require funding and service provision
- research the relationship between socio-economic disadvantage and various health and educational outcomes.³

Prevalence

- In 2017-18, 13.6% of the population or over 1 in 8 (3.24 million people) were estimated to be living below the poverty line, after taking account of their housing costs.⁵
- In 2017-18, there were 489,000 Australian low-income households with children aged 0 -14 years (24% of all low-income households).⁶
- In 2017-18, 17.7% of all Australian children under the age of 15 or over 1 in 6 (774,000 children) were living below the poverty line.⁵
- In 2017-18, in low-income households with children, the average real equivalised disposable income was \$558 per week.⁶
- In 2019, around 11% (289,000) of households with children aged 0 -14 had no paid employment.⁶
- In 2014, 16% of 0 -14-year-old children lived in households deprived of at least 2 essential items - this increased to more than a third (35%) for 1-parent households.⁷
- In 2014, 48% of children in jobless families were deprived of 2 or more essential items.⁷

- In 2014, 1 in 16 (6%) children lived in households that could not afford dental treatment when needed.⁷
- In 2017, 24% or nearly 1 in 4 Australians (4.8 million people) experienced some degree of social exclusion.⁸

Health impacts/outcomes for children

Early childhood development lays the foundation for health, wellbeing, and productivity over the lifespan.⁹ Research has shown that early life cognitive and non-cognitive abilities are important for healthy development throughout childhood and later adult life.¹⁰ Early disadvantage can have harmful effects on children’s development, with more adverse outcomes with each additional factor of disadvantage.⁹ However, not all children have equal opportunities to develop these skills.

Outcomes for children are influenced by the wellbeing of families and the conditions in which they live. Income, finance and employment factors can directly and indirectly affect children, by impacting their education, home environment, housing conditions and access to resources.⁶

A family’s low income can also lead to food insecurity and affect a child’s diet and access to medical care. Appropriate housing, heating and clothing provision can also be impacted by low income, as well as the safety of a child’s environment, and of the quality and stability of their care.⁶

A child’s early years are crucial to shaping their life chances. A child’s development is not only influenced by inherited genes, but also by the quality of family environments, and the availability of appropriate experiences.¹

The cycle of disadvantage as represented below, shows that the risk factors contributing to disadvantage, begin in a child’s first years.¹¹




Figure 1: The cycle of disadvantage

A summary of the research evidence regarding the impact of disadvantage on children’s outcomes, is as follows:

Factor	Outcome
Lower childhood socioeconomic position	<ul style="list-style-type: none"> • Greater risk of experiencing Adverse Childhood Events (ACEs).¹² • Lack of access to resources.¹³ • Children from the lowest socioeconomic areas are more likely to be overweight or obese compared to those from the highest socioeconomic areas.¹⁴
Australian children and adolescents in low-income families, with parents and carers with lower educational levels and higher rates of unemployment	<ul style="list-style-type: none"> • Higher rates of mental disorders in the previous 12 months.¹⁵
Stress from ongoing experiences of disadvantage	<ul style="list-style-type: none"> • Changes brain structure and function, thereby impacting on development.⁹
Children in different socioeconomic groups in Australia	<ul style="list-style-type: none"> • Substantial differences in developmental outcomes on Australian Early Development Census (AEDC) findings.¹⁶ • Widening of gap on all AEDC domains between the percentage of developmentally vulnerable children in the most disadvantaged areas, relative to the least disadvantaged areas from 2009 to 2015. • In 2018 AEDC, this gap is decreasing in the physical health and wellbeing, social competence and emotional maturity domains.¹⁶
<p>Low family income</p> <p>Socioeconomic disadvantage in childhood</p> <p>Children who have experienced poverty at some points in their life</p>	<ul style="list-style-type: none"> • Associated with lower levels of stimulation in the home impacting on children's behaviour and academic skills in early and middle childhood.¹⁷ • Associated with lower working memory ability in children.¹³ • Are likely to have poorer cognitive and social outcomes, are more likely to be obese and are

	also likely to have lower levels of general health. ¹⁸
Children from jobless families	<ul style="list-style-type: none"> • have poorer cognitive and social-emotional outcomes compared to children in families working full-time/long part-time hours.¹

Protective factors

Several studies have found that some protective factors are lessened with experiences of severe neighbourhood disadvantage.¹⁹ However, there is strong evidence to show that education, employment and good health are associated with protection against disadvantage.¹

Protective factors for children in times of adverse experiences include:

- household stability
- a strong early attachment to an adult
- informal support/supportive role model (such as grandparents)
- good parenting.¹

There is also a strong association of improved social and behavioural outcomes for children living in disadvantage with a higher IQ, supportive parenting, and positive parent - child relationships.¹⁹

References
1. McLachlan R, Gilfillan, G. and Gordon, J. Deep and Persistent Disadvantage in Australia. Canberra: Productivity Commission Staff Working Paper; 2013.
2. Australian Bureau of Statistics. 2071.0 - Census of Population and Housing: Reflecting Australia - Stories from the Census, 2016 ABS; 2018 [Available from: https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/2071.0~2016~Main%20Features~Socio-Economic%20Advantage%20and%20Disadvantage~123].
3. Australian Bureau of Statistics. Socio-Economic Indexes for Areas 2018 [Available from: https://www.abs.gov.au/websitedbs/censushome.nsf/home/seifa].
4. Price-Robertson R. What is community disadvantage? Understanding the issues, overcoming the problem: Australian Institute of Family Studies; 2011 [Available from: https://aifs.gov.au/cfca/publications/what-community-disadvantage-understanding-issues-ov].
5. Davidson P, Saunders P, Bradbury B, Wong M. Poverty in Australia: Part 1, Overview. Sydney: ACOSS/UNSW Poverty and Inequality Partnership; 2020.
6. Australian Institute of Health and Welfare. Australia's Children. Canberra: AIHW; 2020.
7. Australian Institute of Health and Welfare. Australia's Children: Material Deprivation Canberra: AIHW; 2022 [Available from: https://www.aihw.gov.au/reports/children-youth/australias-children/contents/income-finance-and-employment-snapshots/material-deprivation].
8. Australian Institute of Health and Welfare. Social determinants of health Canberra: AIHW; 2020 [Available from: https://www.aihw.gov.au/reports/australias-health/social-determinants-of-health].

9. Goldfeld S, O'Connor M, Chong S, Gray S, O'Connor E, Woolfenden S, et al. The impact of multidimensional disadvantage over childhood on developmental outcomes in Australia. *Int J Epidemiol*. 2018;47(5):1485-96.
10. Collier LR, Gregory T, Harman-Smith Y, Gialamas A, Brinkman SA. Inequalities in child development at school entry: A repeated cross-sectional analysis of the Australian Early Development Census 2009–2018. *The Lancet Regional Health - Western Pacific*. 2020;4:100057.
11. The Smith Family. *Breaking the Cycle of Disadvantage*. The Smith Family. 2010.
12. Walsh D, McCartney G, Smith M, Armour G. Relationship between childhood socioeconomic position and adverse childhood experiences (ACEs): a systematic review. *J Epidemiol Community Health*. 2019;73(12):1087-93.
13. Mooney KE, Prady SL, Barker MM, Pickett KE, Waterman AH. The association between socioeconomic disadvantage and children's working memory abilities: A systematic review and meta-analysis. *PloS one* [Internet]. 2021 2021; 16(12):[e0260788 p.].
14. Australian Institute of Health and Welfare. *Childhood Overweight and Obesity—the impact of the home environment*. Canberra: AIHW; 2021.
15. Lawrence D, Johnson S, Hafekost J, Boterhoven De Haan K, Sawyer M, Ainley J, et al. *The Mental Health of Children and Adolescents. Report on the Second Australian Child and Adolescent Survey of Mental Health and Wellbeing*. Canberra: Department of Health; 2015.
16. Commonwealth of Australia. *Australian Early Development Census National Report 2018: A Snapshot of Early Childhood Development in Australia*. Canberra: Department of Education and Training; 2019.
17. Votruba-Drzal E. Economic disparities in middle childhood development: does income matter? *Dev Psychol*. 2006;42(6):1154-67.
18. Warren D. *Low-Income and Poverty Dynamics: Implications for Child Outcomes*. Canberra: Department of Social Services; 2017.
19. Vanderbilt-Adriance E, Shaw DS. Protective factors and the development of resilience in the context of neighborhood disadvantage. *J Abnorm Child Psychol*. 2008;36(6):887-901.

Appendix E: Family domestic violence or conflict

Key points

- Violence is a gendered issue impacting females and children more than males
- Exposure to and experience of FDV impacts health outcomes in children
- Children impacted by FDV have poorer health outcomes than the general child population

Definition

The term 'family and domestic violence' is used in Australia to encompass acts including intimate partner violence, abuse between siblings and other family members and between extended kinship ties.¹

Family and domestic violence is not only physical- it can also include emotional, financial, sexual, verbal, psychological, as well as neglect, coercive control, and stalking.²

Children and young people can experience violence directly (by having violence perpetrated against them) and/or indirectly via witnessing violence being perpetrated against their parent/ or caregiver.²

Prevalence

Family and domestic violence is a major health and welfare issue in Australia and although it does affect people of all ages and from all backgrounds it predominately affects women and children.¹⁻³

It is reported that in Australia one in three women have experienced physical violence and almost one in five women sexual violence. One in four women have experienced physical or sexual violence from their current or former male partner. These figures are most likely an underrepresentation of the actual physical and sexual violence that occurs against women, as only a small proportion of women ever report the violence.²

Children are also victims of violence, either directly or indirectly. Around one in four women report when they experienced violence during a relationship, that children in their care were present and were exposed to the violence, either hearing or seeing the violence.^{2, 3}

There is limited data available on the prevalence and impact of family and domestic and sexual violence on those groups most likely to experience to FDV in Australia. However, the following groups have been identified to be at higher risk of experience and exposure to FDV and consequently higher risk of adverse health and social health outcomes:^{2, 3}

- young women
- women with a lower socio-economic background
- people with disabilities
- children (witness to and experience of FDV)
- Aboriginal people
- people from CALD backgrounds
- LGBTIQ+ people
- people in rural and remote areas

The 2016 COAG report² (pg. 16), noted “the impact of violence against women on some groups of children and young people can be exacerbated by other challenges, including marginalisation and discrimination. This is particularly the case for children who identify as lesbian, gay, bisexual, transgender, intersex and queer or those with parents who identify as such, children from culturally and linguistically diverse backgrounds, and children living in regional, rural and remote areas”.

Health impacts and outcomes- children

Children impacted by FDV are more likely to have poorer health than the general population of children¹ and are more likely to be exposed to other conditions that put them at risk for negative health outcomes.⁴

Children can experience violence as a witness and/or victim.^{2,3} Both these direct and indirect experiences of violence can have long term impact on the child and on the mother-child relationship. Children who are witness to FDV experience similar levels of negative psychological and social issues as those children who are impacted directly by physical abuse.²

Violence against women and their children is the leading cause of homelessness in Australia.^{2,3} In Australia during 2017-18, 22 per cent of clients seeking homelessness services as a result of FDV were aged 0-9 years.³

The impact on children can occur throughout childhood and later in life.^{2,3,5,6} If the violence is chronic or repeated, the symptoms may be exacerbated.²

The exposure of family violence can affect all aspects of a child’s health and wellbeing outcomes,² including during the perinatal period for those children whose mothers experience domestic violence during pregnancy.¹ It affects a child’s mental and physical wellbeing, and can contribute to behavioural issues and poorer educational outcomes.²

FDV does not predetermine outcomes for children and young people; but it can influence them significantly especially when the exposure to the violence occurs in the in early years.⁷

Table 1- Impacts of family and domestic violence on children (adapted from Department for Child Protection⁷ Fact Sheet)

Age of child	Impacts of family and domestic violence
Unborn child	<ul style="list-style-type: none"> • Increased risk of miscarriage, low birth weight and premature birth, foetal injury and death^{2,7} • Weaken developing brains, having lifelong effects on a child's learning, behaviour and health²
Babies and toddlers	<ul style="list-style-type: none"> • Often cry more, show signs of anxiety or irritability • Feeding and sleeping issues • Underweight for age • Neglect • Sexual abuse • Delayed mobility • Often react to loud noises & wary of new people • May be very demanding or very passive • Increased risk of physical injury if in arms of mother whilst assault occurs
Pre-schoolers	<ul style="list-style-type: none"> • Bedwetting, nightmares, eating issues and trouble sleeping • Behavioural issues² such as aggression, lack of emotional control, limited tolerance • Concentration issues • Increased arousal • Physical complaints, fearfulness and numbing • Adjustment problems (i.e. transitioning from kindergarten to pre-primary)
School-age & pre-adolescent	<ul style="list-style-type: none"> • Withdrawal and avoidance from friends and family • Self-harm • Loss of interest in social activities • School performance affected negatively
Adolescents	<p>Increased risk of:</p> <ul style="list-style-type: none"> • academic failure, dropping out of school • delinquency/offending • eating disorders • substance misuse • depression, suicide ideation • use of controlling behaviours • early pregnancy • violent behaviours and violence toward a parent (particularly their mother)

Further impacts of family and domestic violence on children include:

- increased risk of experiencing other forms of abuse such as emotional, physical, or sexual
- higher rates of gastrointestinal problems¹
- higher rates of psychological health issues¹
- increased mental health hospitalisations¹
- hospitalisation from injuries due to abuse including assault, malnutrition and neglect³
- increase risk of homelessness^{2,3}
- increased risk of poor mental health outcomes, particularly depression, anxiety and alcohol dependence^{1,6}
- increased risk of experiencing interpersonal violence as an adult (for both perpetration and victimisation)^{2,6}

Protective factors

Although not well researched, there are attributes or conditions that can occur at an individual, family or community level (protective factors) that can moderate risk or adversity and promote healthy development and child and family wellbeing with regards to exposure and impact on FDV.^{7,8} Not all children are adversely or affected in the same way as a result of exposure to FDV and it is important to consider how children have coped with the violence thus far, what skills and understanding they have developed and what resilience factors have assisted their coping.⁷

Protective factors that may reduce the incidence and severity of impact of FDV (includes child abuse and neglect)⁸ on children include can be grouped into three categories and include:

Individual/child factors

- Social and emotional competence
- Attachment to parent/s

Family/parental factors

- Strong parent/child relationship
- Parental self esteem
- Level of parental education

Social/environmental factors

- Positive social connection and support
- Employment
- Access to health and social services

References

1. Orr C, Fisher CM, Preen DB, et al. Exposure to family and domestic violence is associated with increased childhood hospitalisations. *PLOS ONE* 2020;15:e0237251
2. Commonwealth of Australia, Department of the Prime Minister and Cabinet. COAG Advisory Panel on Reducing Violence against Women and their Children – Final Report. 2016
3. Australian Institute of Health and Welfare. Family, domestic and sexual violence in Australia: continuing the national story 2019. Canberra: AIHW; 2019
4. Development Services Group Inc., Child Welfare Information Gateway. Promoting protective factors for children exposed to domestic violence: A guide for practitioners. Washington, DC Department of Health and Human Services, Administration on Children, Youth and Families, Children's Bureau; 2015
5. Campo M. Children's exposure to domestic and family violence: Key issues and responses (CFCA Paper No. 36). Melbourne: Child Family Community Australia information exchange, Australian Institute of Family Studies; 2015
6. On ML, Ayre J, Webster K, et al. Examination of the health outcomes of intimate partner violence against women: State of knowledge paper. 2016
7. Department of Communities. Fact Sheet 7 Impacts of family and domestic violence on children. In: Child Protection and Family Support, ed. Perth: Government of Western Australia; 2015
8. Child Family Community Australia. Risk and protective factors for child abuse and neglect. CFCA Resource Sheet - May 2017. In: Australian Institute of Family Studies, ed.: Australian Government; 2017

Appendix F: Family Mental Health

Key Points

- Mental health concerns and mental illness are prevalent in Australia
- Mental health can be impacted without having a diagnosis of a mental illness
- Children of parent/s with mental illness are at risk of negative health and developmental outcomes.

Definition

Mental health is an essential component of overall health and wellbeing.¹ The World Health Organization defines mental health as

‘a state of well-being in which every individual realises his or her own potential, can cope with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to her or his community.’²

A mental illness, on the other hand, is a generic term that refers to a group of illnesses. It can be defined as:

‘a clinically diagnosable disorder that significantly interferes with a person’s cognitive, emotional or social abilities’.³

The term mental disorder is also used. Mental illness/disorders include a range of conditions such as:

- anxiety disorders
- affective disorders (e.g. depression)
- psychotic disorders (e.g. schizophrenia), and
- substance use disorders.⁴

It is important to note that a person’s thinking, feeling and behaviour can be impacted by their mental health without meeting the criteria for a mental illness/disorder.⁴

Likewise, it’s possible to be feeling well in many aspects of life while diagnosed with a mental illness²

Many factors both affect and are affected by a person’s mental health. These include their access to services, living conditions and employment status.⁴

Prevalence

Mental Health concerns affect individuals of all ages and backgrounds. The following table describes mental health prevalence data in Australia:

Persons	Prevalence
Mothers of children aged 24 months or less	<p>In 2010:⁵</p> <ul style="list-style-type: none"> • 1 in 5 were diagnosed with depression • Over 50% of those diagnosed, reported that their diagnosed depression was perinatal (that is, the depression was diagnosed from pregnancy until the child was 12 months old). • Of all the cases of diagnosed depression, just over 20% were diagnosed for the first time during the perinatal period.⁵
Children and adolescents	<p>In 2013-14:⁶</p> <ul style="list-style-type: none"> • In the 12 months before the Child and Adolescent Survey of Mental Health and Wellbeing, it is estimated that: <ul style="list-style-type: none"> ◦ 560,000 children and adolescents aged 4 - 17 (14%) experienced a mental health disorder ◦ males had a higher prevalence of mental health disorders (16%) than females (12%) ◦ attention deficit hyperactivity disorder (ADHD) (7.4%); anxiety (6.9%); major depression (2.8%); and conduct disorder (2.1%) were the most prevalent disorders reported by participants.⁶
Children with parents with a mental illness	<ul style="list-style-type: none"> • According to population estimates, 23.3% of all children lived in a family with a parent with a non-substance mental illness.⁷ • 20.4% of mental health service users have dependent children.⁷
Australians aged 16 - 85 years	<ul style="list-style-type: none"> • In 2007, an estimated 1 in 5 (20%) people experienced a mental health issue in the previous 12 months.⁸ • It is expected that almost half (45%) of people in this age group will experience a mental health issue at some point in their life.⁹ • An estimated 2-3% of the population have a severe mental illness (including psychotic disorders and those living with severe depression and anxiety).⁹
People with mental health conditions	<ul style="list-style-type: none"> • In 2019, those with mental health conditions were more likely to drink alcohol at risky levels than those without mental health conditions (21% compared with 17.1% for lifetime risky drinking, and 31% compared with 25% for single occasion risky drinking at least monthly).¹⁰

Some Australians are more likely to experience mental health problems than others. These include:

- young people
- single parent families

- those who are unemployed, and
- Aboriginal people.¹¹

Health impacts/outcomes for child

Mental health is complex. Mental illness can cause distress, and impact on functioning and relationships. It is also associated with poor physical health and early death from suicide.² Mental illness impacts not only the individuals affected, but also those who are around them, including immediate family/children.

The family unit is pivotal for children’s development.¹² It is widely recognised that parental mental health difficulties can impact on children’s development.¹³ Consistent evidence has shown an association between mother’s mental health and children’s adjustment and behaviour.¹⁴ Exposure to adversity at a young age is an established preventable risk factor for mental disorders.¹

The risk of mental illness from parents to children may arise through a complex interplay of risk factors - genetics, neurobiological, as well as a range of psychosocial risk factors - directly by a parent’s behaviour, thoughts and emotions, or indirectly through multiple stressors (such as conflict, isolation, and poverty).¹⁵

A more detailed snapshot of the evidence on children’s outcomes is as follows:

Factor	Outcome/Impact
Children with parent/s who has a mental illness	<ul style="list-style-type: none"> • Have a higher risk of having negative mental health outcomes compared to children of parents without a mental illness.¹⁶ • Have a higher rate of behavioural, developmental, and emotional problems compared to children with parents without a mental illness.^{12, 16} • Are at risk of a similar mental health disorder as their parents and are at risk of a disorders that are specifically related to the parents' diagnosis.¹⁵ • Core attachment needs (such as love, physical and emotional nurturing, and security) may be at risk.¹⁷ • Parenting skills may be impaired including the quality of care and parent-child interaction, with the risk of neglect and potential abuse.¹² • Children may need to assume caring responsibilities for a parent and/or siblings, impacting on age-appropriate activities or school attendance.¹⁸
Adverse Childhood Experiences (ACEs)	<ul style="list-style-type: none"> • Number of ACEs a child is exposed to, is strongly related to the chances of physical and mental health, and social and behavioural problems occurring through childhood into adult life.²⁰

References

1. World Health Organization. Mental health action plan 2013 - 2020. Geneva: WHO; 2013.
2. Beyond Blue. What is Mental Health: Beyond Blue; 2022 [Available from: <https://www.beyondblue.org.au/the-facts/what-is-mental-health>].
3. Council of Australian Governments Health Council. The Fifth National Mental Health and Suicide Prevention Plan. Canberra: Department of Health; 2017.
4. Australian Institute of Health and Welfare. Mental Health Canberra: AIHW; 2020 [Available from: Mental Health].
5. Australian Institute of Health and Welfare. 2010 Australian National Infant Feeding Survey: indicator results. Canberra: AIHW; 2011.
6. Lawrence D, Johnson S, Hafekost J, Boterhoven De Haan K, Sawyer M, Ainley J, et al. The Mental Health of Children and Adolescents. Report on the Second Australian Child and Adolescent Survey of Mental Health and Wellbeing. Canberra: Department of Health; 2015.
7. Maybery D, Reupert A, Patrick K, Goodyear M, L. C. Prevalence of parental mental illness in Australian families. *Psychiatric Bulletin*. 2009;33:22-6.
8. Australian Bureau of Statistics. National Survey of Mental health and Wellbeing: a summary of results 2007. Canberra: ABS; 2008.
9. Department of Health and Ageing. National Mental Health Report 2013: tracking progress of mental health reform in Australia 1993 – 2011. Canberra: Commonwealth of Australia; 2013.
10. Australian Institute of Health and Welfare. Alcohol, tobacco & other drugs in Australia: People with mental health conditions. Canberra: AIHW; 2020.
11. Productivity Commission. Mental Health. Canberra: Commonwealth of Australia; 2020.
12. Reedt C, Lauritzen C, Stover YV, Freili JL, Rognmo K. Identification of Children of Parents With Mental Illness: A Necessity to Provide Relevant Support. *Front Psychiatry*. 2018;9:728.
13. Mensah FK, Kiernan KE. Parents' mental health and children's cognitive and social development: families in England in the Millennium Cohort Study. *Soc Psychiatry Psychiatr Epidemiol*. 2010;45(11):1023-35.
14. Giallo R, Cooklin A, Wade C, D'Esposito F, M. N. Maternal postnatal mental health and later emotional-behavioural development of children: the mediating role of parenting behaviour. *Child: Care Health and Development*. 2014;40(3):327-36.
15. van Santvoort F, Hosman CM, Janssens JM, van Doesum KT, Reupert A, van Loon LM. The Impact of Various Parental Mental Disorders on Children's Diagnoses: A Systematic Review. *Clin Child Fam Psychol Rev*. 2015;18(4):281-99.
16. Drost LM, van der Krieke L, Sytema S, Schippers GM. Self-expressed strengths and resources of children of parents with a mental illness: A systematic review. *Int J Ment Health Nurs*. 2016;25(2):102-15.
17. Robinson E, Rodgers B, P. B. Family relationships and mental illness: impacts and service responses. In: Australian Institute of Family Studies, editor.: Australian Family Relationships Clearinghouse; 2008.
18. Reupert A, Maybery D, M. K. Children whose parents have a mental illness: prevalence, need and treatment. *Medical Journal of Australia*. 2013;199(3):57-9.
19. Emerging Minds. Adverse Childhood Experiences (ACEs): Summary of evidence and impacts [Available from: <https://emergingminds.com.au/resources/adverse-childhood-experiences-aces-summary-of-evidence-and-impacts/>].
20. Hughes K, Bellis MA, Hardcastle KA, Sethi D, Butchart A, Mikton C, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health*. 2017;2(8):e356-e66.

21. Curran E, Adamson G, Stringer M, M. R. Severity of mental illness as a result of multiple childhood adversities: US National Epidemiologic Survey. *Social Psychiatry and Psychiatric Epidemiology*. 2016;51:647-57.
22. Liu SR, Kia-Keating M, Nylund-Gibson K, Barnett ML. Co-Occurring Youth Profiles of Adverse Childhood Experiences and Protective Factors: Associations with Health, Resilience, and Racial Disparities. *Am J Community Psychol*. 2020;65(1-2):173-86.
23. Turney K. Maternal depression and childhood health inequalities. *Journal of Health and Social Behavior*. 2011;52(3):314-32.
24. Clavarino AM, Mamun AA, O'Callaghan M, Aird R, Bor W, O'Callaghan F, et al. Maternal anxiety and attention problems in children at 5 and 14 years. *J Atten Disord*. 2010;13(6):658-67.
25. Foster K, Lewis P, McCloughen A. Experiences of peer support for children and adolescents whose parents and siblings have mental illness. *J Child Adolesc Psychiatr Nurs*. 2014;27(2):61-7.

Appendix G: Homelessness, transience and/or overcrowding and remoteness

Key Points

- Homelessness, overcrowding and housing stress are widespread issues impacting children, young people and their families across Western Australia
- Homelessness, overcrowding and housing stress have both immediate and longer term impacts on a child's development and their physical, mental, emotional and social health.
- There are specific protective factors that can reduce the negative impacts on a child of homelessness, overcrowding and housing stress.
- Clinical staff need to be sensitive to the health, wellbeing and developmental impacts of homelessness, overcrowding and housing stress when planning and delivering care for this cohort.

Definition

Homelessness

There are many accepted definitions of homelessness.

In alignment with the Western Australian government's 10-Year strategy¹ on homelessness, the Australian Bureau of Statistics (ABS) definition of homelessness has been adopted here. The ABS define a person as homeless if they do not have suitable accommodation alternatives and their current living arrangement:

- is in a dwelling that is inadequate,
- has no tenure, or if their initial tenure is short and not extendable,
- or does not allow them to have control of, and access to space for social relations.²

Homelessness Australia, in addition to the ABS definition, consider a cultural definition of homelessness which includes three categories of homelessness:³

- Primary homelessness – is experienced by people without conventional accommodation (e.g. those who are rough sleeping)
- Secondary homelessness – is experienced by people who frequently move from one temporary shelter to another (e.g. those who are 'couch surfing')
- Tertiary homelessness – is experienced by people staying in accommodation that falls below minimum community standards.

Overcrowding

Overcrowding occurs when a dwelling is too small for the size and composition of the household living in it. The Australian Government uses the Canadian National Occupancy Standard (CNOS) as a measure of overcrowding. CNOS states that an overcrowded dwelling is one that requires at least 1 additional bedroom.⁴ The CNOS measure specifies that: ⁴

- No more than 2 people share a bedroom
- Parents or couples may share a bedroom
- Children under 5, either of the same sex or opposite sex may share a bedroom

- Children under 18 of the same sex may share a bedroom
- A child aged 5-17 should not share a bedroom with a child under 5 of the opposite sex
- Single adults 18 and over and any unpaired children require a separate bedroom.

Housing Stress

In general housing stress is experienced when housing costs are high relative to income. In these situations, the costs of housing are likely to reduce the household's ability to afford other living costs such as food, clothing, transport and utilities.⁵ A household living with housing stress is defined as one that spends more than 30% of their gross income on housing costs.⁵

Prevalence

Homelessness

Some children live in families experiencing homelessness, whilst others experience homelessness on their own.⁶ Poverty is a key driver of homelessness for children and their families. Homelessness for children and young people occurs, most commonly, through the following pathways:

- being part of a homeless family,
- leaving the family home with one parent (usually to escape violence or abuse),
- leaving the family home independently,
- exiting the care or youth justice system.⁷

On Census night 2016 there were almost 2000 Western Australian children experiencing homelessness.² This figure is known to be an underestimate due to the hidden nature of homelessness.

Australian children are more likely to experience homelessness if they:

- live in remote and very remote areas,
- live in multiple family households (compared to those living in single parent or couple family households),
- live in areas of greater socioeconomic disadvantage,
- identify as Aboriginal or Torres Strait Islander.⁶

In 2020-21, 7102 children aged 0 to 17 years presented at WA specialist homelessness services alone or with their families.⁸ The majority of these children were aged under 10 years.⁹ The single most common reason for children and young people to need housing and homelessness assistance is family and domestic violence.⁹ While the proportion of male and female young people aged 10 to 14 years presenting to homelessness services in WA was relatively even, for young people aged 15 to 17 years the proportion of female clients was greater than that of male clients (61.7% female compared to 38.3% male).⁷ Almost 50% of all young people aged 10 to 17 years presenting to homelessness services are Aboriginal.⁸

Overcrowding

On Census night 2016 approximately 18,900 children aged 0 to 14 years were living in overcrowded housing.² In June 2020, it was estimated that 8% of households were in overcrowded dwellings.⁴ One-quarter of state government owned and managed Aboriginal households were in overcrowded dwellings across Australia.⁴ Children living in low socioeconomic areas were 12 times more likely to be living in an overcrowded dwelling as those from high socioeconomic areas.¹⁰

Housing stress

In WA, 21.3% of WA children aged 0 to 14 years live in a household experiencing housing stress.⁵ Almost half of WA single parents who live in rented accommodation experience housing stress, with 40% of WA single parents who are homeowners also experience housing stress.¹¹ Housing stress is more common in major cities than in remote areas.¹⁰

Health and developmental impacts/outcomes for children

Homelessness

Homelessness and housing stress have both immediate and longer-term effects on a child's health and wellbeing.⁷ Experiences of homelessness affect physical health, educational attainment and social functioning.⁷ Families and children experiencing homelessness are likely to experience social exclusion, compromised safety, and lack of connectedness with the school and broader community.¹²

Preschool and school-aged children experiencing homelessness are more likely to experience:

- Mental health problems
- Emotional or behavioural problems and
- Food insecurity (which could potentially lead to adverse physical health).⁵

The overall youth unemployment rate in January 2022 was approximately 9%, with the unemployment rate for adults sitting at 4% for the same period.¹³ Unemployment in people who have experienced homelessness as a child is much higher. Children who first experience homelessness under 15 years of age have an employment rate of just 10% by the time they're adults, as opposed to 24% if they're homeless after 15 years.¹² Children experiencing homelessness are also at increased risk of being homeless as adolescents and adults.¹⁴

Overcrowding

Those living in overcrowded housing may not be able to access basic amenities which are necessary for health, including washing, laundry, hygienic food storage and preparation, and safe disposal of waste.¹⁴ Overcrowding can increase the risk of family conflict or violence, child abuse and neglect.¹⁵ Additionally, overcrowding has been associated with increased risk of emotional and behavioural problems and reduced school performance.¹⁰

Overcrowding is higher among Aboriginal households. This can have a detrimental effect on Aboriginal children's ear and skin health.¹⁰

Housing Stress

Children in households not experiencing housing stress often have better health and school engagement.¹⁰ Housing stress can:

- Negatively impact on parental mental health
- Reduce investment in children's food, health and education
- Increase a child's risk of material deprivation and social exclusion.¹⁰

Protective Factors

The Center on the Developing Child at Harvard University identifies four factors that can lead to positive outcomes when a child is facing adversity;¹⁶

- The opportunity for the child to improve their adaptive and self-regulatory skills
- Facilitating supportive adult-child relationships
- Spiritual connections and cultural traditions
- Building a sense of self-efficacy and perceived control

There are protective factors that can reduce the negative impacts of homelessness in children and their families. Some of these include:¹⁴

- Staying with others, such as a partner, friends or family (though not necessarily in accommodation)
- Having activities that they enjoy and having these activities planned
- Having a pet.

Protective factors specifically for young people experiencing homelessness, include:¹⁷

- Having a connection to an adult
- Having at least a high school education
- Being currently enrolled in school or having a full-time job.

For Aboriginal children and their families, a strong connection to Aboriginal culture is a strength and protective factor against adversity. The values of kinship, interdependence, group cohesion and community loyalty are protective factors for Aboriginal families against homelessness.¹

References

1. Department of Communities. All Paths Lead to a Home: Western Australia's 10-Year Strategy on Homelessness 2020-2030. Perth, Western Australia: Western Australian Government; 2020
2. Australian Bureau of Statistics. Census of Population and Housing: Estimating Homelessness. Canberra, Australia; 2018
3. Homelessness Australia. About Homelessness. 2022. Available at: <https://homelessnessaustralia.org.au/about-homelessness/>. Accessed 4th April, 2022
4. Australian Institute of Health and Welfare. Housing assistance in Australia Canberra, Australia: Australian Government; 2021

5. Australian Institute of Health and Welfare. Children's Headline Indicators: Housing Stress. Canberra, Australia: Australian Government; 2018
6. Australian Institute of Health and Welfare. Australia's children: Homelessness. Canberra, Australia: Australian Government; 2022
7. Commissioner for Children and Young People Western Australia. Indicators of Wellbeing: Experiences of homelessness and housing stress. 2022. Available
8. Commissioner for Children and Young People Western Australia. Profile of Children and Young People in WA - 2022. Perth: Commissioner for Children and Young People Western Australia; 2022
9. Australian Institute of Health and Welfare. Specialist Homelessness Services Collection (SHSC) data cubes 2011-12 to 2020-21. Canberra, Australia: Australian Government; 2021
10. Australian Institute of Health and Welfare. Australia's children. Canberra, Australia: Australian Government; 2022
11. Bankwest Curtin Economics Centre. Getting our House in Order? BCEC Housing Affordability Report 2019. Focus on Western Australia Report Series. Perth: Curtin University; 2019
12. Emerging Minds. Families and homelessness: Supporting parents and improving outcomes for children. Webinar: Australian Institute of Family Studies; 2021
13. Australian Bureau of Statistics. Labour Force, Australia. 2022. Available. Accessed 14th April, 2022
14. Flatau P, Tyson K, Callis Z, et al. The State of Homelessness in Australia's Cities: A Health and Social Cost Too High. Perth, Western Australia: Centre for Social Impact The University of Western Australia.; 2018:2-3
15. Cant RL, O'Donnell M, Sims S, et al. Overcrowded housing: One of a constellation of vulnerabilities for child sexual abuse. *Child Abuse and Neglect* 2019;93:239-248
16. Centre on the Developing Child Harvard University. Key Concepts - Resilience. 2022. Available at: <https://developingchild.harvard.edu/science/key-concepts/resilience/>
17. Kelly P. Risk and protective factors contributing to homelessness among foster care youth: An analysis of the National Youth in Transition database. *Children and Youth Services Review* 2020;108
18. Kaleveld L, Seivwright A, Flatau P, et al. Ending Homelessness in Western Australia 2019 Report. The Western Australian Alliance to End Homelessness Annual Snapshot Report Series. Perth: The University of Western Australia, Centre for Social Impact.; 2019

Appendix H: Trauma

Key points

- Trauma is an experience/s that induces an abnormally intense and prolonged stress response. For children it is often the result of adverse childhood experiences (ACEs)
- Trauma experiences can affect a child's physical, social, emotional and behavioural wellbeing
- Nurses need to be sensitive to the potential presence of trauma history when working with children and young people and use a trauma informed care approach to practice

Definitions

Trauma is an experience/s that induces an abnormally intense and prolonged stress response. It often involves a sense of intense fear, terror, and helplessness. It involves experiences that overwhelms a person's ability to cope.¹

Complex trauma

"Occurs when a child repeatedly experiences severe stressors of traumatic events over an extended period of time".^{2,3} These stressors start at a developmental time point in in childhood when the child is considered vulnerable. Examples of complex trauma include:

- Physical, social or sexual abuse
- Neglect
- Witnessing family and domestic violence

Complex trauma results from more than one traumatic event that disrupts the development of emotional health and the regulation of emotions.² It affects the ability to have clear thoughts or memories and can disrupt a child's sense of safety and trust in important relationships.²

Intergenerational trauma

"Is the impact of trauma experienced in parents'/family/caregivers lives being passed down to their children. Intergenerational trauma is often discussed in the context of Aboriginal and Torres Strait Islander young people, and among children of refugees. It can also be experienced by children of veteran and other parents continuing to be affected by their own trauma".⁴

Adverse childhood experiences (ACE)

Perceived negative events that are outside the control of the child, which may hinder normal development, cause harm or the potential for harm, and are accompanied by stress and suffering.⁵ It is a term used to describe very stressful events or circumstances that children may experience in their childhood.

Prevalence

Although the total prevalence of childhood trauma is difficult to measure, it is common. The literature suggests between 57-75 per cent of Australians will experience a potentially traumatic event during their lifetime.⁴ The consequences of early trauma can continue through a person's life.⁶

Trauma exposure is more common among specific groups including:^{4, 7}

- people who experience homelessness
- young people in out-of-home care or under youth justice supervision
- refugees
- women and young people experiencing family and domestic violence
- LGBTIQ young people
- Aboriginal and Torres Strait Islander young people

Trauma is more common in women and in population groups that have a higher incidence of cumulative or intergenerational trauma (for example refugee or Aboriginal communities).⁶ For Aboriginal and Torres Strait Islander young people, the trauma experiences are more likely to be complex with both the impact of intergenerational and current trauma experiences.⁴

Trauma is often a result of adverse childhood experiences (ACEs).^{5, 8, 9} ACEs include harms that affect children directly and indirectly through their living environment:

- family difficulties related to financial struggles,
- family health problems and losses,
- parenting impairment due to mental health condition or alcohol and other drug misuse,
- marital discord and parental and family conflicts,
- physical, emotional and sexual abuse,
- neglect- physical and emotional.

ACE's can have a long-lasting negative effect on a person's health; however poor outcomes are not inevitable. Not every child is negatively affected by trauma and adversity; the presence and reinforcement of protective factors can help in developing resilience and reducing the negative impact of trauma.⁵

Health impacts/outcomes for child

Trauma experiences can affect a child's social, emotional and behavioural wellbeing. Repeated trauma increases vulnerability in children.² Exposure to ACEs during childhood and adolescence can result in significant developmental delays, lower educational attainment and social and emotional instability.⁵

Disruptions in attachment

Infancy is a crucial time in the development of attachment relationships and is the most vulnerable to disruptions, with complex trauma compromising the development of a secure child-caregiver attachment relationship.^{2, 3} Possible indicators include heightened discriminate attachment behaviour, with insecure, anxious or disorganised attachment behaviour.

If a child has experienced disruptive attachment in their early years, they may struggle understanding and forming subsequent relationships. It may also have a flow on effect for the child's social and emotional development.

In early childhood, difficulties in attachment may present as:¹⁰

- clingy, difficulty with separations
- inconsistent behaviour towards caregiver/s
- lacking in trust towards others
- finding it challenging to seek help from others
- struggling to self-regulate emotions and behaviours

Changes to child's stress response

When there is a real or perceived threat a child's stress response system is activated to prepare them to 'fight' or 'flee'. When an 'infant or young child experiences trauma events, their body's stress system may be excessively and repeatedly activated'.^{2, 10} Additionally, the development and regulation of the body's stress response can be disrupted -this can mean that the reactions to stress can be blunted or exaggerated.

Changes to brain development

During early childhood development, brain development is particularly sensitive to trauma. Trauma events may disrupt brain structure and function. Changes to brain areas may contribute to many symptoms of complex trauma including:^{2-4, 10}

- social, emotional and behavioural difficulties
- developmental issues such as speech, language and cognitive difficulties

Trauma and mental ill-health

A history of trauma that has impacted on development is the single most significant predictor for the presentation to mental health services as an adult.¹¹ Individual trauma results from an event, series of events or set of circumstances that is experienced by an individual as physically or emotionally harmful or life threatening and has lasting adverse effects on the individual's functioning and mental, physical, social, emotional or spiritual well-being.¹²

Trauma exposure can:^{3, 4}

- increase the risk of mental ill-health
- lengthen the duration of the illness
- compound its severity and complexity and
- impact on treatment response.

Trauma related mental health diagnosis can include 'PTSD, anxiety, depression, psychosis, personality disorders, self-harm and suicide related behaviours, eating disorders and comorbidity with alcohol and substance misuse'.⁴

Social and emotional wellbeing and functioning

Social wellbeing and functioning can be affected by trauma experiences in childhood, impacting on a child's ability to form relationships and friendships. In young children these social struggles may present as:²

- difficulty trusting others and feeling safe in a relationship
- feelings of fear, threat, rejection or being unloved when socialising
- vigilance or guardedness when interacting with others
- struggle to interact with authority figures, such as educators
- struggle with social skills

Emotional wellbeing and functioning are also impacted by trauma, with many people who have experienced complex trauma struggling to regulate their emotions. This can lead to people living with strong feelings of shame, self-blame and low self-esteem.³

Impact of Intergenerational Trauma on child development

The *Make Healing Happen*¹³ report states 'adverse experiences in childhood can have lifelong effects. Traumatic childhood experiences, such as those of Stolen Generations survivors, may affect following generations through biological changes in stress responses and by undermining the ability to parent and love freely without fear'.

These traumatic experiences can be transferred from the first generation of survivors that have experienced (or witnessed) it directly in the past to the second and further generations of descendants of the survivors.¹⁴

This can create developmental issues for children, who are particularly susceptible to distress at a young age. These children may experience difficulties with attachment, disconnection from their extended families and culture, and high levels of stress from family and community members who are dealing with the impacts of trauma. It can also create a cycle of trauma, where the impact is passed from one generation to the next.^{14, 15}

ACEs and impact on health outcomes

There is relationship between multiple ACEs and the increased risk of poor health outcomes across various health related behaviours and conditions.⁹ There are varying associations between multiple ACEs (4 or more) and;⁹

- vulnerability to mental health conditions in childhood, adolescence and adult life,^{5, 6, 9} which can impair social and emotional functioning.^{5, 7, 16}
- increased and problematic alcohol and drug use
- sexual risk taking
- self-directed violence
- physical inactivity, obesity and diabetes
- cancer, heart disease and respiratory disease

The evidence on the harms that multiple ACEs place on health throughout the life-course highlights the importance of addressing the various stressors that can occur in a child's life.⁹ A reduction in ACEs and building resilience to enable those affected, to avoid their harmful effects could have a major effect on health outcomes.⁹

Protective factors

There are protective factors (both for parents/caregiver and child) that if present and reinforced in a child's life, can build resilience and reduce the impacts of adversity.

These include:

Child

- Safe, caring and supportive relationship with someone they trust^{5, 16}
 - research has shown that having one positive caring relationship can substantially improve a child's recovery and healing from trauma
 - having someone that makes the child feel safe and protected helps support their mental health and resilience

- Promoting and developing a child's resilience⁹
- Social and emotional competence of children¹⁷

Parent

- Parental resilience
- Knowledge of parenting and child development¹⁷
- Social connections¹⁷
- Concrete supports for children¹⁷

The Center on the Developing Child at Harvard University identifies four factors that can lead to positive outcomes when a child is facing adversity;¹⁶

- The opportunity for the child to improve their adaptive and self-regulatory skills
- Facilitating supportive adult-child relationships
- Spiritual connections and cultural traditions
- Building a sense of self-efficacy and perceived control

References

1. headspace. The impact of trauma on mental health. 2022. Available at: <https://headspace.org.au/explore-topics/for-young-people/trauma/>. Accessed 14/04/2022,
2. Michele Hervatin, Emerging Minds. National Workforce Centre for Child Mental Health. Complex trauma through a trauma informed lens: Supporting the wellbeing of infants and young children. Australia: Australian Institute of Family Studies; 2021
3. Blue Knot Foundation. Understanding Trauma and Abuse. Available at: <https://blueknot.org.au/resources/understanding-trauma-and-abuse/what-is-childhood-trauma/>. Accessed 28/01/2022, 2022
4. Bendall S, Phelps A, Browne V, et al. Trauma and young people. Moving toward trauma-informed services and systems Melbourne: Orygen, The National Centre of Excellence in Youth Mental Health; 2018
5. Emerging Minds. National Workforce Centre for Child Mental Health, Australian National University. Adverse Childhood Experiences (ACEs): Summary of evidence and impacts. 2020
6. Wei-May Su, Louise Stone. Adult survivors of childhood trauma: Complex trauma, complex needs. *Australian journal of general practice* 2020;49:423-430
7. Australian Institute of Health and Welfare. Stress and Trauma. 2020. Available at: <https://www.aihw.gov.au/reports/australias-health/stress-and-trauma>. Accessed 09 February 2022, 2022
8. Bartlett JD, Smith S. The role of early care and education in addressing early childhood trauma. *American Journal of Community Psychology* 2019;64:359-372
9. Hughes K, Bellis MA, Hardcastle KA, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *The Lancet Public Health* 2017;2:e356-e366

10. Georgetown University Centre for Child and Human Development (GUCCHD). Recognising and addressing trauma in infants, young children, and their families Washington, DC: Georgetown University.; 2020
11. Kaiser S, Zimmet M, Fraser J, et al. Recognition of attachment difficulties and developmental trauma is the responsibility of all paediatricians. *J Paediatr Child Health* 2018;54:1110-1116
12. Substance Abuse Mental Health Service Administration. SAMHSA's concept of trauma and guidance for a trauma-informed approach. 2014
13. Healing Foundation. Make Healing Happen. It's time to act. Canberra; 2021
14. Atkinson J. Trauma-informed services and trauma specific care for Indigenous Australian children. Resource sheet no. 21 produced for the Closing the Gap Clearinghouse. In: Australian Institute of Family Studies, Australian Institute of Health and Welfare, eds. Canberra; 2013
15. SNAICC- National Voice for our Children. Family Matters Report: 2021.; 2021
16. Centre on the Developing Child. Harvard University. Key Concepts - Resilience. 2022. Available at: <https://developingchild.harvard.edu/science/key-concepts/resilience/>. Accessed 21 March, 2022
17. Child Welfare Information Gateway. Protective factors approaches in child welfare. In: U.S. Department of Health and Human Services. Children's Bureau, ed. Washington, D.C.; 2020